

# Intronic hammerhead ribozymes are ultraconserved in the human genome

Marcos de la Peña<sup>+</sup> & Inmaculada García-Robles

Instituto de Biología Molecular y Celular de Plantas, Universidad Politécnica de Valencia-Consejo Superior de Investigaciones Científicas, Valencia, Spain

**Small ribozymes have been regarded as living fossils of a prebiotic RNA world that would have remained in the genomes of modern organisms. In this study, we report the ultraconserved occurrence of hammerhead ribozymes in Amniota genomes (reptiles, birds and mammals, including humans), similar to those described previously in amphibians and platyhelminth parasites. The ribozymes mapped to intronic regions of different genes, such as the tumour suppressor RECK in birds and mammals, a mammalian tumour antigen and the dystrobrevin beta in lizards and birds. *In vitro* characterization confirmed a high self-cleavage activity, whereas analysis of RECK-expressed sequence tags revealed fusion events between the *in vivo* self-cleaved intron and U5 or U6 small nuclear RNA fragments. Together, these results suggest a conserved role for these ribozymes in messenger RNA biogenesis.**

**Keywords:** ribozyme; intron; splicing; exaptation; retrotransposons

EMBO reports (2010) 11, 711–716. doi:10.1038/embor.2010.100

## INTRODUCTION

The term ‘RNA world’ was first used to refer to a hypothetical stage in the emergence of life in which RNA carried out both genetic and catalytic roles (Gilbert, 1986). This hypothesis stimulated research not only on the origin of life but also on new RNA capabilities, ultimately unveiling new features of our ‘modern RNA world’. The discovery of catalytic RNAs in the Group I intron (Cech *et al*, 1981) and in RNase P (Guerrier-Takada *et al*, 1983), the more recent findings of RNA interference (Fire *et al*, 1998) and the ribozymal nature of peptide bond formation (Nissen *et al*, 2000) are all hallmarks of the ever-growing biological roles of RNA. In this context, previous data have revealed the presence of small self-cleaving RNA domains in the noncoding regions of mammalian genes, such as the co-transcriptional cleavage ribozyme at the 3′-untranslated region of

$\beta$ -globin (Teixeira *et al*, 2004), an intronic Hepatitis  $\delta$  virus-like ribozyme (HDVR) in *Cpeb3* (Salehi-Ashtiani *et al*, 2006), and a discontinuous hammerhead ribozyme (HHR) at the 3′-untranslated region of *Clec2* genes (Martick *et al*, 2008).

In this study, we show the results of a genome-wide search for the HHR domain, a paradigmatic ribozyme initially described to catalyse a transesterification self-cleavage reaction during replication of some viroids and other small RNA plant pathogens (Prody *et al*, 1986). Since then, HHRs have also been observed in a few unrelated eukaryotic genomes, such as those of salamanders (Epstein & Gall, 1987), schistosomes (Ferbeyre *et al*, 1998), crickets (Rojas *et al*, 2000), *Arabidopsis thaliana* (Przybilski *et al*, 2005) and *Clec2* genes of rodents and platypus (Martick *et al*, 2008). In the latter case, HHRs showed a clear similarity to those observed in schistosomes, suggesting an event of horizontal gene transfer (Martick *et al*, 2008). On the basis of this similarity, we followed a bioinformatic search among lower metazoan and vertebrate genomes that revealed hundreds of HHR motifs associated with retrotransposable elements. More surprisingly, a few similar ribozymes were also detected as ultraconserved motifs in the genomes of Amniota spp., indicating ancient exaptation events for this small self-cleaving domain during the evolution of higher vertebrates.

## RESULTS


### An iterative method for identifying HHR-like motifs

Extensive bioinformatic approaches have been devised previously to search for HHR motifs among genomes (Ferbeyre *et al*, 1998, 2000; Przybilski *et al*, 2005; Martick *et al*, 2008), exclusively considering the 11 catalytically conserved nucleotides as the source of the phylogenetic signal (Fig 1). In this study, we used an iterative bioinformatic approach starting with basic local alignment search tool (BLAST) searches among vertebrate genomes, followed by the implementation of structural and phylogenetic constraints (supplementary information online). Briefly, initial queries (seeds) were composed by naturally occurring Helix II motifs (supplementary Fig S1 online) flanked by the conserved U and P boxes (Fig 1A). Sequence changes in the returned entries were examined to fulfil two conditions: (i) not to affect the conserved U and P boxes and (ii) to be either compensatory within stem II or located in loop 2. Filtered hits were analysed for three

Instituto de Biología Molecular y Celular de Plantas, Universidad Politécnica de Valencia-Consejo Superior de Investigaciones Científicas, Avenida de los Naranjos s/n, 46022 Valencia, Spain

<sup>+</sup>Corresponding author. Tel: +34 963877861; Fax: +34 963877859; E-mail: [rivero@ibmcp.upv.es](mailto:rivero@ibmcp.upv.es)

Received 18 March 2010; revised 11 May 2010; accepted 10 June 2010; published online 23 July 2010



**Fig 1** | Sequence strings of 22–32 nucleotides were used for searching HHRs. (A) Schematic representation of the seeds used to search for HHRs. Conserved motifs corresponding to the U and P boxes are shown in black squares. (B) Schematic representation of type I/II (left) and II/III (right) HHRs. Helix II, U and P boxes are depicted in black and white. Consensus self-cleavage site (RUH box), Helix I and Helix III, not included in the seeds, are depicted in grey. HHR, hammerhead ribozyme.

additional criteria to define candidate sequences as HHR motifs: (iii) 5' and 3' surrounding regions should fold as two helices (I and III), wherein (iv) Helix I had to be either a loop-closed RNA helix (type I/II HHR) or an open helix containing an internal loop (type II/III HHR), and (v) a proper cleavage site triplet must exist (Fig 1B). An extra point of validation was the presence and nature of loops 1 and 2, which allow the required tertiary interactions for *in vivo* self-cleavage activity (de la Peña *et al*, 2003, 2009; Khvorova *et al*, 2003; Martick & Scott, 2006).

### HHRs are widespread in amphibians and lampreys

The BLAST searches using the *Schistosoma mansoni* genome (Berriman *et al*, 2009) revealed the presence of more than 50,000 entries for type II/III HHRs (data not shown). Ribozymes were similar to those described previously in the SM $\alpha$  satellite DNA of these platyhelminth parasites (Fig 2A; Ferbeyre *et al*, 1998), although their occurrence was not restricted to repetitive DNA. Similar HHRs were also detected in the genome of *Schistosoma japonicum* (Fig 2B), and in a few complementary DNA (cDNA) sequences of the more distant liver fluke *Opisthorchis viverrini* (Fig 2C). Interestingly, one of these cDNAs showed significant homology to an endonuclease reverse transcriptase from *S. japonicum* (supplementary Table S1 online).


Similar type II/III HHRs embedded in 330bp tandem repeats (satellite 2 DNA) have been described previously in the genomes of several species of newts (Epstein & Gall, 1987) and salamanders (Caudata order; Zhang & Epstein, 1996). Our iterative searches revealed new HHRs in the genomes of three more species of salamanders and especially in the genome of the frog *Xenopus tropicalis* (Anura order), suggesting a widespread occurrence among amphibians (Fig 2D–E; supplementary Table S1 online). BLAST-like alignment tool (BLAT; University of California, Santa Cruz) searches using whole HHRs of *X. tropicalis* revealed hundreds of entries dispersed along the frog genome. Some of

these ribozymes were observed to be associated with repetitive DNA, but they were also dispersed as single intronic motifs noticeably associated with reverse transcriptase-like coding regions (supplementary Table S1 online).

Type II/III HHRs were also detected in the genomes of *Petromyzon marinus* (marine lamprey) and *Lethenteron japonicum* (arctic lamprey; Fig 2F). BLAT searches further confirmed the presence of hundreds of single HHRs dispersed along the *P. marinus* genome, which in many cases also mapped within intronic regions (supplementary Table S1 online). Although a detailed analysis of HHRs observed in schistosomes, *X. tropicalis* and *P. marinus* will be reported elsewhere (M.D.P., unpublished observations), their occurrence in different intronic regions and their frequent association with reverse transcriptase-like genes was striking.

### A HHR is ultraconserved in a tumour suppressor gene

Parallel BLAST searches against the human genome were performed with Helix II seeds derived from HHRs of schistosomes and lower vertebrates. For several seeds (supplementary information online), a hit mapping within a contig of the human chromosome 9 was repeatedly obtained (Fig 3A). Surrounding regions of this hit could be folded to give an HHR fold (hereafter referred to as HH9). The HH9 fold showed a 4 bp Helix III capped by a non-palindromic tri-loop and a 13 bp Helix I containing a 7 nt internal loop (Fig 3B). *In vitro* characterization of HH9 revealed a  $k_{\text{obs}}$  of  $2.43 \pm 0.22 \text{ min}^{-1}$  at 25 °C, pH 6.5 and 1 mM  $\text{Mg}^{2+}$  (supplementary Fig S2 online), which corresponds to a self-cleavage activity similar to those described for most naturally occurring HHRs acting *in cis* (de la Peña *et al*, 2003; Khvorova *et al*, 2003). As reported for the discontinuous HHR in *Clec2* genes (Martick *et al*, 2008), loop 2 of HH9 and also loop 1 (not included in the searching seeds) were similar to the corresponding loops of the *S. mansoni* HHRs, allowing equivalent loop–loop tertiary interactions (supplementary Fig S2C online; Martick & Scott, 2006). Human HH9 mapped to the middle of the sixth intron (14kb) of an open reading frame coding for the protein RECK, a factor characterized initially as a tumour suppressor (Takahashi *et al*, 1998). This membrane glycoprotein is known to inhibit the matrix metalloproteinases involved in remodelling the extracellular matrix, a key step during embryogenesis and vasculogenesis (Oh *et al*, 2001). The HH9 fold was observed to be ultraconserved (90% identity for 60 nt) among the introns of RECK orthologues for all endothermic vertebrate spp. examined (birds and mammals), but was absent in the orthologues of lower vertebrates and metazoans (Fig 3A; supplementary Fig S3 online). Furthermore, sequence heterogeneity was mostly restrained to the Helix III–loop 3 region and was always compatible with a catalytically active HHR. Sequence alignment of the surrounding intronic regions of HH9 did not show any significant conservation, therefore revealing a strong selection pressure in the ribozyme motif (Fig 3B; supplementary Fig S3 online). Data mining of RECK expressed sequence tags revealed two examples from *Bos taurus* fetal and calf brain tissues, which mapped exactly to the precise 5' self-cleaved HH9. However, both sequences showed 5' extensions of 82 and 83 nt that corresponded to fragments of U5 and U6 small nuclear RNAs (snRNAs), respectively (Fig 3C). These two examples reveal that HH9 self-cleaves *in vivo*, and also offers circumstantial evidence that the 5'-OH product could interfere with the spliceosomal machinery.



**Fig 2** | HHR motifs in trematodes and lower vertebrates. (A) Consensus example of the HHR with some of the natural variability described previously for three schistosomal spp. (Ferbeyre *et al*, 1998). Examples of HHRs detected in (B) *Schistosoma japonicum* (variability with respect to *Schistosoma mansoni* is highlighted) and (C) *Opisthorchis viverrini* (intraspecies variability is highlighted). Examples of HHRs observed in (D) *Caudata* (salamanders and newts), (E) *Anura* (frog) and (F) *Agnatha* (lampreys) spp. HHR, hammerhead ribozyme.

### A HHR in a mammalian tumour antigen gene

By using the iterative search strategy described above, a second HHR motif (HH10) was detected in human chromosome 10. The HH10 motif was similar to HH9, with the exception of a larger Helix III (43 nt instead of 11 nt; Fig 4A), and was conserved among most mammalian genomes from platypus to humans, with the exception of rodents (rat and mouse), lagomorphs (rabbit) and cingulata (armadillo; supplementary Fig S4 online). In humans, the motif mapped within the first intron (10 kb) of the uncharacterized *C10orf118* gene. The protein encoded by this open reading frame is the CTCL (for cutaneous T-cell lymphoma) tumour antigen L14-2, a factor involved in processes of tumorigenesis (Hartmann *et al*, 2004). The surrounding regions of HH10 Helix I revealed a clear conservation of eight extra base pairs when compared with the HH9 region (Figs 3A, 4A; supplementary Fig S4 online). Furthermore, the 5'-ends of two expressed sequence tags (both from human brain origin) were observed to match with this intronic region. However, the 5' extremes did not map with the precise self-cleavage site, but with the end of the conserved region of Helix I (supplementary Fig S5 online). Regarding the absence of the HH10 motif among some mammal species, a case of an inactive HHR motif was observed in the *C10orf118* orthologue of the lagomorph *Ochotona princeps* (supplementary Fig S6 online), which suggests the loss of this HHR motif during the evolution of these mammals. Interestingly, a putative gene homologous to *C10orf118* (UniProt accession number C5J917) was detected in the *S. mansoni* genome. Inspection of the corresponding genomic sequence in the trematode revealed the presence of at least two HHR motifs (supplementary Fig S7 online).


### A conserved HHR in the DTNB gene of sauropsids


The BLAT searches using the whole HH9 motif revealed a new example of a conserved HHR (HHsau) in genomes of four sauropsids: the *Anolis carolinensis* lizard and *Gallus gallus*, *Taenopygia guttata* and *Meleagris gallopavo* birds. The structure and sequence of HHsau mostly resembled HH9, although a covariation within stem I matched with the one detected in HH10 (Fig 4B). For the four spp., HHsau mapped within the first intron (9 kb) of a gene encoding the dystrobrevin beta protein (DTNB). DTNB is expressed in brain tissues, where it interacts with dystrophin to form the core of the transmembrane dystrophin glycoprotein complex (DGC; Peters *et al*, 1997). Although DGC is known to link the extracellular matrix to the intracellular cytoskeleton, the specific presence of DTNB involves the DGC in neuronal signalling (Grady *et al*, 2006). The mammalian orthologues of DTNB seem to lack the extra first exon/intron present in sauropsids, which would explain the absence of this particular HHR in mammals.

### DISCUSSION

In this study, we have unveiled the presence of ultraconserved intronic HHRs in the genomes of reptiles, birds and mammals, including humans. To achieve this goal, we first identified a large set of HHRs in the genomes of trematodes, *P. marinus* and *X. tropicalis* through iterative bioinformatic searches. The recurrent association of many of these motifs with RT-like coding regions would directly involve these ribozymes in retrotransposable elements, extending previous observations of HHR in the satellite DNA of newts, trematodes and crickets to a more general landscape among eukaryotes and other organisms (this work; de la Peña &







**Fig 4** | Intronic HHRs conserved in mammals and sauropsids.

(A) Secondary structure of the human HH10. Sequence heterogeneity detected in the genomes of alpaca (Vp), cow (Bt), dog (Cf), dolphin (Tt), elephant (La), flying fox (Pv), guinea pig (Cp), hedgehog (Ee), mouse lemur (Mm), pig (Ss), tarsier (Ts), treeshrew (Tb) and sloth (Ch) is depicted in red. Nucleotides in black are conserved in all mammals having a HH10 motif. Red arrows point to the 5'-ends of human expressed sequence tags mapping at this region. (B) HHRs observed in the first intron of the dystrobrevin beta gene in *Anolis carolinensis* (in green), *Taeniopygia guttata* (in pink), *Gallus gallus* (in red) and *Meleagris gallopavo* (in brown). Positions in black are identical to the human HH9. A covariation similar to the one detected in HH10 is shown in blue. HHR, hammerhead ribozyme; mRNA, messenger RNA; snRNA, small nuclear RNA.

Nevertheless, the occurrence of intronic HHRs in Amniota genomes as ultraconserved motifs within specific genes strongly suggests that they have a conserved biological role other than retrotransposition. Previous studies using artificially engineered HHRs into introns did not reveal splicing effects on adjacent exons in wild-type yeast (Lacadie *et al*, 2006) and human cells (Dye *et al*, 2006). However, more recently it has been shown that continuous precursor messenger RNAs (pre-mRNAs) seem to be required for proper splicing (Fong *et al*, 2009). In that work, the authors observed that, although highly active HDVR promotes splicing inhibition, an intronic HHR affects splicing weakly, most probably because of its lower catalytic efficiency. Accordingly, we propose that natural intronic HHRs might not affect pre-mRNA splicing under basal conditions, but either changes in transcription elongation or ribozyme upregulation could interfere with the spliceosomal machinery (for example, fusion with snRNAs), promoting alternative splicing events or even disruption of the pre-mRNA processing. A similar situation is illustrated by the intronic HDVR of the mammalian *Cpeb3* gene, which has been described already as a low-activity version of this ribozyme (Salehi-Ashtiani *et al*, 2006). Altogether, these findings allow us to conclude that self-cleaving ribozymes present in retrotransposons of lower metazoans would have been exapted or domesticated in the genomes of Amniotes to perform specific functions.

Finally, the resemblance of vertebrate HHRs with those widespread within mobile genetic elements in trematodes suggests that recurrent genomic horizontal transfers could have taken place from parasites to hosts. In fact, increasing evidence for a relationship between carcinogenesis and infections by helminths, specifically trematodes, has accumulated during the last decades (WHO, 1994; Mayer & Fried, 2007), and our present data adds circumstantial support in this direction. Although a deeper molecular characterization is required, a potential relationship can be envisaged between trematode-induced carcinogenesis and ribozyme-containing mobile elements, which would ultimately affect the host genomic stability. In a more unlikely scenario, trematode infection could promote activation of the intronic HHRs of host genes involved in tumour progression.

## METHODS

**Analysis of data.** Genome servers used for the analysis of data and mapping of hits were NCBI-BLAST2 Nucleotide tool at the European Bioinformatics Institute (www.ebi.ac.uk), ENSEMBL

García-Robles, 2010). Furthermore, recent work describing the association of the HDVR motif with retrotransposons (Webb *et al*, 2009) confirms that not only HHR, but self-cleaving ribozymes in general might function in retrotransposition.

(www.ensembl.org), University of California Santa Cruz Genome Bioinformatics (www.genome.ucsc.edu) and the National Center for Biotechnology Information (www.ncbi.nlm.nih.gov).

**In vitro transcription.** *Cis*-acting HHRs were synthesized by *in vitro* transcription of linearized plasmids containing the corresponding cDNA inserts immediately preceded and followed by the promoter of T7 RNA polymerase and the *Xba*I site, respectively. Transcription reactions were carried out in the presence of P32-labelled UTP and blocking deoxyoligonucleotides to avoid extensive self-cleavage. Resulting primary transcripts were fractionated by polyacrylamide gel electrophoresis (PAGE) in 15% gels with 8 M urea and eluted.

**Cis cleavage kinetics under protein-free conditions.** Uncleaved primary transcripts were incubated in 20 µl of 50 mM PIPES-NaOH (pH 6.5) for 1 min at 95 °C and slowly cooled to 25 °C for 15 min. After taking a zero-time aliquot, self-cleavage reactions were triggered by adding 1 mM MgCl<sub>2</sub>. Aliquots removed at appropriate time intervals were separated by PAGE in 15% denaturing gels. The product fraction at different times,  $F_t$ , was determined by quantitative scanning and fitted to the equation  $F_t = F_0 + F_\infty(1 - e^{-kt})$ , where  $F_0$  and  $F_\infty$  are the product fractions at zero time and at the reaction end point, respectively, and  $k$  is the first order rate constant of cleavage ( $k_{obs}$ ).

**Supplementary information** is available at *EMBO reports* online (<http://www.emboreports.org>).

#### ACKNOWLEDGEMENTS

We thank W.G. Scott, L. Yenush, S. Delgado and J. de la Peña for advice and critical reading of the paper. This work was supported by Ministerio de Educación y Ciencia of Spain (BFU2008-03154/BMC and Ramón y Cajal contract to M.D.P.) and Generalitat Valenciana (GV06/206 to M.D.P.).

#### CONFLICT OF INTEREST

The authors declare to have a patent pending (no. P201030569) at the Spanish Patent and Trademark Office (OEPM, Madrid, Spain).

#### REFERENCES

- Berriman M *et al* (2009) The genome of the blood fluke *Schistosoma mansoni*. *Nature* **460**: 352–358
- Cech TH, Zaug AJ, Grabowski PJ (1981) *In vitro* splicing of the ribosomal RNA precursor of tetrahymena: involvement of a guanosine nucleotide in the excision of the intervening sequence. *Cell* **27**: 487–496
- de la Peña M, Gago S, Flores R (2003) Peripheral regions of natural hammerhead ribozymes greatly increase their self-cleavage activity. *EMBO J* **22**: 5561–5570
- de la Peña M, Dufour D, Gallego J (2009) Three-way RNA junctions with remote tertiary contacts: a recurrent and highly versatile fold. *RNA* **15**: 1949–1964
- de la Peña M, García-Robles I (2010) Ubiquitous presence of the hammerhead ribozyme motif along the tree of life. *RNA* (in press)
- Dye MJ, Gromak N, Proudfoot NJ (2006) Exon tethering in transcription by RNA polymerase II. *Mol Cell* **21**: 849–859
- Epstein LM, Gall JG (1987) Self-cleaving transcripts of satellite DNA from the newt. *Cell* **48**: 535–543
- Ferbeyre G, Smith JM, Cedergren R (1998) Schistosome satellite DNA encodes active hammerhead ribozymes. *Mol Cell Biol* **18**: 3880–3888
- Ferbeyre G, Bourdeau V, Pageau M, Miramontes P, Cedergren R (2000) Distribution of hammerhead and hammerhead-like RNA motifs through the GenBank. *Genome Res* **10**: 1011–1019
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **391**: 806–811
- Fong N, Ohman M, Bentley DL (2009) Fast ribozyme cleavage releases transcripts from RNA polymerase II and aborts co-transcriptional pre-mRNA processing. *Nat Struct Mol Biol* **16**: 916–922
- Gilbert W (1986) Origin of life: the RNA world. *Nature* **319**: 618
- Grady RM, Wozniak DF, Ohlemiller KK, Sanes JR (2006) Cerebellar synaptic defects and abnormal motor behavior in mice lacking alpha- and beta-dystrobrevin. *J Neurosci* **26**: 2841–2851
- Guerrier-Takada C, Gardiner K, Marsh T, Pace N, Altman S (1983) The RNA moiety of ribonuclease P is the catalytic subunit of the enzyme. *Cell* **35**: 849–857
- Hartmann TB, Thiel D, Dummer R, Schadendorf D, Eichmüller S (2004) SEREX identification of new tumour-associated antigens in cutaneous T-cell lymphoma. *Br J Dermatol* **150**: 252–258
- Khvorova A, Lescoute A, Westhof E, Jayasena SD (2003) Sequence elements outside the hammerhead ribozyme catalytic core enable intracellular activity. *Nat Struct Biol* **10**: 708–712
- Lacadie SA, Tardiff DF, Kadener S, Rosbash M (2006) *In vivo* commitment to yeast cotranscriptional splicing is sensitive to transcription elongation mutants. *Genes Dev* **20**: 2055–2066
- Martick M, Scott WG (2006) Tertiary contacts distant from the active site prime a ribozyme for catalysis. *Cell* **126**: 309–320
- Martick M, Horan LH, Noller HF, Scott WG (2008) A discontinuous hammerhead ribozyme embedded in a mammalian messenger RNA. *Nature* **454**: 899–902
- Mayer DA, Fried B (2007) The role of helminth infections in carcinogenesis. *Adv Parasitol* **65**: 239–296
- Nissen P, Hansen J, Ban N, Moore PB, Steitz TA (2000) The structural basis of ribosome activity in peptide bond synthesis. *Science* **289**: 920–930
- Oh J *et al* (2001) The membrane-anchored MMP inhibitor RECK is a key regulator of extracellular matrix integrity and angiogenesis. *Cell* **107**: 789–800
- Peters MF, O'Brien KF, Sadoulet-Puccio HM, Kunkel LM, Adams ME, Froehner SC (1997) Beta-dystrobrevin, a new member of the dystrophin family. Identification, cloning, and protein associations. *J Biol Chem* **272**: 31561–31569
- Prody GA, Bakos JT, Buzayan JM, Schneider IR, Bruening G (1986) Autolytic processing of dimeric plant virus satellite RNA. *Science* **231**: 1577–1580
- Przybilski R, Gräf S, Lescoute A, Nellen W, Westhof E, Steger G, Hammann C (2005) Functional hammerhead ribozymes naturally encoded in the genome of *Arabidopsis thaliana*. *Plant Cell* **17**: 1877–1885
- Rojas AA, Vazquez-Tello A, Ferbeyre G, Venanzetti F, Bachmann L, Paquin B, Sbordoni V, Cedergren R (2000) Hammerhead-mediated processing of satellite pDo500 family transcripts from *Dolichopoda* cave crickets. *Nucleic Acids Res* **28**: 4037–4043
- Salehi-Ashtiani K, Lupták A, Litovchick A, Szostak JW (2006) A genomewide search for ribozymes reveals an HDV-like sequence in the human *CPEB3* gene. *Science* **313**: 1788–1792
- Takahashi C *et al* (1998) Regulation of matrix metalloproteinase-9 and inhibition of tumor invasion by the membrane-anchored glycoprotein RECK. *Proc Natl Acad Sci USA* **95**: 13221–13226
- Teixeira A, Tahiri-Alaoui A, West S, Thomas B, Ramadass A, Martianov I, Dye M, James W, Proudfoot NJ, Akoulitchiev A (2004) Autocatalytic RNA cleavage in the human beta-globin pre-mRNA promotes transcription termination. *Nature* **432**: 526–530
- Webb CT, Riccitelli NJ, Ruminski DJ, Lupták A (2009) Widespread occurrence of self-cleaving ribozymes. *Science* **326**: 953
- WHO (1994) Schistosomes, liver flukes and *Helicobacter pylori*. IARC working group on the evaluation of carcinogenic risks to humans. *IARC Monogr Eval Carcinog Risks Hum* **61**: 1–241
- Zhang Y, Epstein LM (1996) Cloning and characterization of extended hammerheads from a diverse set of caudate amphibians. *Gene* **172**: 183–190